

2.2 Taking samples

Using a stainless steel knife, roughly 1 kg of edible portions of various imported and local vegetables were obtained. In November, the following vegetables were picked from Plate 1 of the Erbil market: pepper (*Capsicum annuum*), tomato (*Solanum L.*), and eggplant (*Solanum melongena L.*). To get rid of dust and debris, tap and distilled water were used to clean each vegetable sample. The edible parts of the samples were dried in an oven for a full 72 hours. After the materials were dried, they were finely powdered and put in containers to wait for analysis. Luminescence Spectrometer for X-rays XRF was used to examine the different forms of heavy metals, including lead (Pb), manganese (Mn), zinc (Zn), and cadmium (Cd). The mechanism by which XRF operates involves the displacement of electrons from their atomic orbital positions, resulting in the release of an energy burst that is unique to a given element. The XRF instrument's detector then records this energy release and classifies the energies according to their constituent elements.



Plate 1: Nearby Grocery and Convenience Stores, sample location.

3. RESULTS AND DISCUSSION

The findings showed that all of the samples contained the heavy metals that were found, albeit in different amounts. Table 1 displays the FAO/WHO standard concentrations of heavy metals in vegetables (mg/kg). Table 2 displays the heavy metal concentrations for local tomatoes, peppers, and eggplants; Table 3 displays the concentrations for imported vegetables. Lead is the metal with the highest concentration found in both imported and local eggplant, according to researchers results (Younis and Darwesh's 2023). The least amount of metals found in both imported and local tomatoes and peppers is cadmium; in eggplant, both local and imported zinc exceeded FAO/WHO acceptable limits; in tomato and pepper, both imported and local zinc fell short of FAO/WHO acceptable limits. The current findings concurred with those of researcher results in 2018, who reported zinc concentrations in tomatoes of 17.3 mg kg⁻¹ (Hasoon, 2018). The manganese content of both imported and local tomatoes was high, and the other metal contents of all vegetables were found to be higher than FAO/WHO acceptable limits in both cases Owing to various factors such as The use of specific fertilizers and pesticides, mining operations, and industrial pollution can all lead to the buildup of heavy metals in the soil. Vegetables may take up these metals and store them in their tissues if they are grown in contaminated soil. In a similar vein, a group of researchers discovered that the concentrations of heavy metals in various plant parts varied (Santamaria et al., 1999). If veggies are kept in heavy metal-containing containers. These metals may be present in the materials used to make the containers, or they may have become contaminated during production or storage. Vegetables stored in areas with high air pollution levels may become contaminated due to the presence of heavy metals in the air. In urban areas or close to industrial facilities, this is more likely to happen. Vegetables can get contaminated by cross-contamination if they are kept close to other products that are contaminated with heavy metals, like batteries or electronic gadgets. Vegetables may get contaminated if they are kept in water that has been tainted by heavy metals.

Table 1: FAO/WHO standard concentrations of heavy metals in vegetables (mg/kg).				
Types of vegetables	Pb	Mn	Zn	Cd
Tomato	0.1	10	20	0.1
Eggplant	0.3	3.3	30	0.05
Pepper	0.3	2	50	0.1

Table 2: Heavy metal concentration of four local vegetable samples (mg/kg).				
Types of vegetables	Pb	Mn	Zn	Cd
Tomato	40	15.2	18.3	3
Eggplant	102	30	52	7
Pepper	50.2	12	15	5

Table 3: Heavy metal concentration of four imported vegetable samples (mg/kg).				
Types of vegetables	Pb	Mn	Zn	Cd
Tomato	55	15	16.2	7
Eggplant	122	30	55	8
Pepper	62.3	32	39.7	7

4. CONCLUSION

The only heavy metal with concentrations below the limits set by the Food and Agriculture Organization and the World Health Organization was zinc. Zinc concentrations in tomatoes were lower in both imported and local varieties, similar to the lower concentrations in peppers. However, zinc concentrations in both imported and local eggplant were higher. The levels of lead and cadmium found in both domestic and imported vegetables are higher than those allowed by the FAO/WHO. There could be a number of reasons for this, including: Contamination of soil: Heavy metals from industrial processes like smelting and mining can build up in soil. or from using pesticides and fertilizers that have been tainted. If grown in contaminated soil, vegetables may absorb these metals through their roots and store them in their tissues. Water contamination: Heavy metals can also enter the food chain through water pollution. If vegetables are grown in contaminated areas or are irrigated with tainted water, they may absorb these metals and accumulate them in their tissues. Air pollution: Heavy metals such as lead and mercury have the ability to enter the atmosphere and land on crops. This could occur near busy roads or industrial zones. If heavy metals are present in the machinery or packaging materials used during processing and packaging, vegetables may also become contaminated with heavy metals.

REFERENCES

Akan, J.C., Kolo, B.G., Yikala, B.S. and Ogugbuaja, V.O., 2013. Determination of some heavy metals in vegetable samples from Biu local government area, Borno State, North Eastern Nigeria. *International Journal of Environmental Monitoring and Analysis*, 1 (2), Pp. 40-46.

Akter, S., Goto, A. and Mizoue, T., 2017. Smoking and the risk of type 2 diabetes in Japan: a systematic review and meta-analysis. *Journal of epidemiology*, 27 (12), Pp. 553-561.

Aysha, M.I.J., Zakir, H.M., Haque, R., Quadir, Q.F., Choudhury, T.R., Quraishi, S.B. and Mollah, M.Z.I., 2017. Health risk assessment for population via consumption of vegetables grown in soils artificially contaminated with arsenic. *Archives of Current Research International*, 10 (3), Pp.1-12.

Elbagermi, M.A., Edwards, H.G.M. and Alajtal, A.I., 2012. Monitoring of heavy metal content in fruits and vegetables collected from production and market sites in the Misurata area of Libya. *International Scholarly Research Notices*, 2012.

Hassoon, H.A., 2018. Heavy metals contamination assessment for some imported and local vegetables. *Iraqi Journal of Agricultural Sciences*, 49 (5).

Ihekoronye, A.I. and Ngoddy, P.O., 1985. *Integrated food science and technology for the tropics* (pp. ix+-386pp).

Li, X., Li, Z., Lin, C.J., Bi, X., Liu, J., Feng, X., Zhang, H., Chen, J. and Wu, T., 2018. Health risks of heavy metal exposure through vegetable consumption near a large-scale Pb/Zn smelter in central China. *Ecotoxicology and environmental safety*, 161, Pp. 99-110.

Mahdavi, M., Amin, M.M., Mahvi, A.H., Pourzamani, H. and Ebrahimi, A., 2018. Metals, heavy metals and microorganism removal from spent filter backwash water by hybrid coagulation-UF processes. *Journal of Water Reuse and Desalination*, 8 (2), Pp.225-233.

Mahmood, A., Mahmoud, A.H., El-Abedein, A.I.Z., Ashraf, A. and Almunqedhi, B.M., 2020. A comparative study of metals concentration

- in agricultural soil and vegetables irrigated by wastewater and tube well water. *Journal of King Saud University-Science*, 32 (3), Pp. 1861-1864.
- Santamaria, P., Elia, A., Serio, F. and Todaro, E., 1999. A survey of nitrate and oxalate content in fresh vegetables. *Journal of the Science of Food and Agriculture*, 79 (13), Pp.1882-1888.
- Solidum, J., Dykimching, E., Agaceta, C. and Cayco, A., 2012. Assessment and identification of heavy metals in different types of cooked rice available in the Philippine market. In 2nd international conference on environmental and agriculture engineering IPCBEE, 37, Pp. 35-39. Singapore: IACSIT Press.
- Tariq, F.S., Samsuri, A.W., Karam, D.S., Aris, A.Z. and Jamilu, G., 2019. Bioavailability and mobility of arsenic, cadmium, and manganese in gold mine tailings amended with rice husk ash and Fe-coated rice husk ash. *Environmental monitoring and assessment*, 191, Pp.1-12.
- Yang, Q.W., Xu, Y., Liu, S.J., He, J.F. and Long, F.Y., 2011. Concentration and potential health risk of heavy metals in market vegetables in Chongqing, China. *Ecotoxicology and environmental safety*, 74 (6), Pp. 1664-1669.
- Youni, A.M. and Darwesh, D.A., 2023. Assessment of Heavy Metals in Some Local and Imported Vegetables in Erbil Market. *Kirkuk Journal of Science*, 18 (4).
- Yusuf, K.A. and Oluwole, S.O., 2009. Heavy metal (Cu, Zn, Pb) contamination of vegetables in urban city: a case study in Lagos. *Research Journal of Environmental Sciences*, 3 (3), Pp.292-298.

